

Proceedings of the 11th SKAM Lithic Workshop  
The multifaceted biface - Bifacial technology in Prehistory  
20th–22nd of October, 2014, Miskolc, Hungary



## Bifacial technology at the beginning of the Upper Paleolithic in Moravia

Petr Škrdla

<b>Abstract</b>	Leaf points have been documented in the Szeletian, Bohunician and Aurignacian technocomplexes in Moravia. The leaf point undoubtedly represents a Szeletian type-fossil within the Moravian early Szeletian, but in the Bohunician the presence of leaf points is more debatable. Bifacial knapping is an unusual element within the Bohunician Levallois-based industry, and has only been documented in the type-site assemblage. Leaf points in Aurignacian contexts have never been recorded in stratified assemblages; they have been documented only at surface sites so their association with the Aurignacian is doubtful.
<b>Kivonat</b>	<b>Bifaciális technológia Morvaországban a felső paleolitikum kezdete idején</b> Morvaországban (Cseh Köztársaság) a Szeletai, a Bohunicien és az Aurignacien technokomplexekhez egyaránt rendelünk kétoldali megmunkálású levéleszközöket. A levélhegy kétség kívül a morvaországi korai Szeletai vezérellete, a Bohunicienben való jelenléte viszont vitatott. A kétoldali megmunkálás a Levallois-alapú Bohunicienben szokatlan jelenség, csak a névadó lelőhely eszközkészletéből ismert ilyen kidolgozású darab. Aurignacien levéleszközök rétegzett lelőhely készleteinek kontextusában nem ismertek, csupán felszíni gyűjtések során kerültek elő, kapcsolatuk az Aurignacien-nel így kérdéses.
<b>Keywords</b>	<i>Late Middle Palaeolithic, Upper Palaeolithic, Aurignacian, Bohunician, Szeletian, leaf points</i>
<b>Kulcsszavak</b>	<i>Késői középső paleolitikum, felső paleolitikum, Aurignacien, Bohunicien, Szeletai ipar, levéleszközök</i>
<b>Author / Szerző</b>	Academy of Sciences of the Czech Republic, Institute of Archaeology in Brno 602 00 Brno, Czech Republic. E-mail: ps@iabrno.cz
<b>Cite as / Hivatkozás</b>	Škrdla, P. (2016) Bifacial technology at the beginning of the Upper Paleolithic in Moravia. <i>Litikum</i> 4: 5–8.
<b>Article history / Kézirat történet</b>	Received // Érkezés: 2015. 12. 03. Accepted // Elfogadás: 2016. 01. 12. Published // Közzététel: 2016. 04. 29.
<b>Copyright / Jogok</b>	© 2016 Škrdla. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. // Ez egy nyílt hozzáférésű publikáció, amit a Creative Commons 4.0 licensze véd. A termék szabadon használható, terjeszthető és sokszorosítható az eredeti szerző és forrás megjelölése mellett.



### 1. Introduction

The beginning of the Upper Paleolithic in Moravia (Fig. 1) is characterized by two so-called transitional technocomplexes – Szeletian and Bohunician – followed by the Evolved Aurignacian that first appears after Heinrich Event 4. Chronologically, both the Bohunician and the Szeletian appear during Greenland Interstadial (GI) 12 (Škrdla et al. 2014, Fig. 10), although the results of luminescence dating (Richter et al. 2008; Nejman et al. 2011) suggest that they may have already appeared during the preceding interstadial (GI 13).


### 2. Szeletian

The Szeletian in Moravia has been reported from many sites (Valoch 2012), but only three sites (Vedrovice V, Moravský Krumlov IV and Želešice III) have yielded diagnostic assemblages from stratified contexts. The Szeletian industry is characterized by the frequent occurrence of bifacial tools – including leaf points supplemented by different varieties

of side scrapers, points, and end scrapers. Bifacial thinning flakes (BTFs) which represent characteristic byproducts of bifacial thinning (Fig. 2) (Škrdla et al. 2014) present additional evidence for bifacial reduction.

### 3. Bohunician

In contrast to the Szeletian, leaf points and flat retouch do not seem to play an important role in stratified Bohunician assemblages. No leaf points have been found at the Stránská skála site cluster, Tvarožná, and Ořechov IV assemblages. The only exception where the leaf points play a more significant role is the Bohunice type site, where a series of leaf points (Fig. 3), BTFs and flat-retouched tools were recovered. There are also dozens of surface collections (e.g. Líšeň, Mohelno, Ondratice) where leaf points and artifacts produced using the Bohunician Levallois technology have been reported. Surface collections often originate from large sites located on significant positions in the landscape, so they may represent palimpsests resulting from sequential




**Figure 1.** Map showing sites mentioned in text. 1: Bohunice, 2: Boršice / Buchlovice, 3: Lhota / Hlinsko, 4: Líšeň I, 5: Mohelno, 6: Moravský Krumlov IV, 7: Ondratice / Želeč, 8: Ořechov IV, 9: Stránská skála, 10: Tvarožná X, 11: Vedrovice V, 12: Želešice III. //

**1. ábra.** A cikkben említett lelőhelyek. 1: Bohunice, 2: Boršice / Buchlovice, 3: Lhota / Hlinsko, 4: Líšeň I, 5: Mohelno, 6: Moravský Krumlov IV, 7: Ondratice / Želeč, 8: Ořechov IV, 9: Stránská skála, 10: Tvarožná X, 11: Vedrovice V, 12: Želešice III.

occupations by people with differing technocomplexes. The presence of leaf points in the Bohunice assemblage has been discussed by many authors. All hypotheses have been summarized and tested by Tostevin and Škrdla (2006) using material from the Bohunice 2002 excavation. This collection was excavated using modern methodological standards (including proveniencing of artifacts using a laser total station, wet sieving of sediments) and we were able to reject the “Excavation Bias” hypothesis and Oliva’s (1981, 1984) “Traded Point” hypothesis. However, three other hypotheses and namely the “Pedogenic”, “Sequential Occupation”, and “Landscape” hypotheses, cannot be currently rejected. In order to test the latter three hypotheses, we considered raw material, technological, and typological homogeneity / heterogeneity of Bohunician collections. In recent years, we applied a new survey methodology focused on discovering new stratified sites (Škrdla et al. 2016) and subsequently conducted excavations at Tvarožná, Ořechov, Ondratice/Želeč, and Líšeň where artifacts within intact sediments were discovered. To date, we have obtained diagnostic collections from only two sites – Tvarožná X (6.9 km east of Stránská skála) and Ořechov IV (13.9 km west of Stránská skála). As the leaf points and BTFs are absent in both collections, the “landscape” hypothesis can probably be rejected. This project is ongoing and we are currently investigating site formation processes.

AMS dates suggest a long time-span for the existence of the Bohunician and we can hypothesize two or more chronologically distinct phases - one with and one without leaf points. We also need to take into account both the chronological overlap with the Szeletian and the presence of Stránská skála chert originating directly from the Stránská skála rock outcrop within the Želešice III Szeletian assemblage (Škrdla et al. 2014) – both facts may have resulted in Szeletian influence on Bohunician Levallois-based technology. However, the distribution of dates from site to site currently does not support this patterning. In addition, the problems



**Figure 2.** Bifacial thinning flakes from Želešice III. //  
**2. ábra.** Bifaciális felületalakító szilánkok Želešice III. lelőhelyről.

with reliability of dates from this time period (Jöris, Street 2008) will currently not allow us to test this hypothesis.

#### 4. Aurignacian

Although several surface sites classified as Aurignacian contain leaf points, they have not been documented within any stratified collection, which makes their association with the Aurignacian technocomplex disputable. The so-called industries “Aurignacian with leaf points” including “Morava-type Aurignacian” (Klíma 1978) and “Míšovice-type industry” (Oliva 1990), are based on surface collections which may not be homogeneous. An alternative hypothesis for the presence of bifacially worked artifacts within Aurignacian assemblages asserts that the leaf points in Aurignacian contexts represent prepared bladelet cores (or a lithic form, in which raw material was transported from outcrops in general; cf. Oliva’s (1981) “handaxe” from Karolín), rather than projectile points. This hypothesis is supported by several broken leaf points that possess bladelet removals from the broken surface along their “crested” edge (Fig. 4).


#### 5. Triangular bifacially worked points

Triangular bifacially worked points have a specific character and they are particularly concentrated in the Lhota / Hlinsko site cluster (Moravian Gate) with isolated items also collected at sites outside of this site-cluster (Škrdla 2007). Again, all sites are surface sites. Although those artifacts were collected within assemblages together with characteristic Aurignacian artifacts, they could indicate some influence of the Kostenki-Streletskaya technocomplex from the east (Škrdla 2007).

#### 6. Conclusion

The Szeletian technocomplex is characterized by ubiquitous bifacially worked tools including leaf points of different shapes.

In sites classified as Bohunician, leaf points have been documented only in the Bohunice type-site assemblage, where they are made from a different raw material than the rest of Levallois industry. There are sites Stránská skála (III, IIIa, IIIc, IIa), Tvarožná X and Ořechov IV, where



**Figure 3.** Bohunician leaf points, Bohunice 2002 excavation. //  
**3. ábra.** Bohunician levélhegyek, Bohunice 2002-es ásatás.


bifacially worked artifacts are not present. Bifacial knapping is also not present at some Eurasian IUP sites including Boker Tachtit (Negev Desert, Israel; Marks, Kaufman 1983), Kremenets-Kulychivka (Ternopil Area, Western Ukraine; Škrdla, Nikolajev 2014), and Kara Bom (Altai Republic, Russian Federation; Derevianko, Petrin, Rybin, 2000). An important question that needs to be asked is: Are the leaf points actually an integral part of the Bohunician (or, in general, IUP) technocomplex? Or do they merely represent local innovation / intrusion?

Although several evolved Aurignacian (~30–33 <sup>14</sup>C BP) assemblages have been reported, none of them yielded any leaf points. On the other hand, the bifacial reduction can be interpreted as part of raw material transport, or bladelet core preparation.

Triangular bifacially worked points have never been discovered in a stratified context. The chronological categorization of this technocomplex and its possible relationship to the Kostenki-Streletskaia technocomplex are currently unresolved issues.

### Acknowledgments

The completion of the manuscript was supported by GACR project #15-19170S.



**Figure 4.** Bladelet removals on broken leaf points from the Aurignacian surface site of Boršice / Buchlovice. //

**4. ábra.** Mikropenge-leválasztások nyomai törött levéleszközökön Boršice / Buchlovice felszíni Aurignacien lelőhelyről.

### References

- Derevianko, A.P., Petrin, V.P., Rybin, E.P. 2000. The Kara-Bom Site and the Characteristics of the Middle-Upper Paleolithic Transition in the Altai. *Archaeology, Ethnology and Anthropology of Eurasia* 2 (2): 33–52.
- Jöris, O. Street, M. 2008. At the end of the 14C time scale—the Middle to Upper Paleolithic record of western Eurasia. *Journal of Human Evolution* 55 (5): 782–802.

- Klíma, B. 1978. Paleolitická stanice u Přestavlk, okr. Přerov. *Archeologické rozhledy* 30: 5–13.
- Marks A.E., Kaufman D. 1983. Boker Tachtit: The Artifacts. In: Marks A.E. (ed.), *Prehistory and Paleoenvironments in the Central Negev, Israel. The Avdat/Aqev Area, Part 3*. Dallas, TX: Southern Methodist University Press, 69–126
- Nejman, L., Rhodes, E., Škrdla, P., Tostevin, G., Neruda, P., Nerudová, Z., Valoch, K., Oliva, M., Kaminská, L., Svoboda, J., Grün, R. 2011. Chronological Review of the Middle to Upper Palaeolithic Transition in the Czech Republic and Slovakia: New Optically Stimulated Luminescence results. *Archaeometry* 53 (5): 1044–1066.
- Oliva, M. 1981. Acheulian finds from Karolín, District of Kroměříž (Czechoslovakia). *Anthropologie* 19: 27–32.
- Oliva, M. 1990. La signification des pointes foliacées dans l'Aurignacien et dans le Type de Míškovice. In: Kozłowski, J. K. (ed.), *Les industries à pointes foliacées du Paléolithique supérieur Européen*. ERAUL 42, Liege, 224–232.
- Richter, D., Tostevin, G. & Škrdla, P. 2008. Bohunician technology and thermoluminescence dating of the type locality of Brno-Bohunice (Czech Republic). *Journal of Human Evolution* 55 (5): 871–885.
- Škrdla, P. 2007. Paleolitické osídlení Hlinska a okolí. In: Šebela, L. a kol., *Hlinsko. Výšinná osada lidu badenské kultury*. Spisy Archeologického ústavu AÚ AV ČR Brno 32. Brno, AÚ AV ČR, 31–47.
- Škrdla, P., Nejman, L., Rychtaříková, T., Nikolajev, P., Lisá, L. 2014. New observations concerning the Szeletian in Moravia. *Quartär* 61: 87–101.
- Škrdla, P., Nejman, L., Rychtaříková, T. 2016: Early Upper Paleolithic Sites in Southern Moravia: A Method for Finding Stratified Sites. *Journal of Field Archaeology* 41(1): 57–67.
- Tostevin G.B., Škrdla P. 2006. New excavations at Bohunice and the question of the uniqueness of the type-site for the Bohunician industrial type. *Anthropologie* 44: 31–48.
- Valoch, K. 2012. Szeletien na Moravě a na západním Slovensku. *Acta Musei Moraviae, Scientiae sociales* 97 (2): 167–198.

## SKAM 2014 Article

This study was presented at the 11th SKAM Lithic Workshop: the multifaceted biface - Bifacial technology in Prehistory. 20th–22nd of October, 2014, Miskolc, Hungary.

The conference papers are published in the Litikum Journal volumes as special contributions. Informations about the conference as well as the abstract book are available on the SKAM 2014 website: <http://skam.pannontenger.hu>